

Calidad de Vida: a systematic review of quality of life in Latino cancer survivors in the USA

Cleo A. Samuel^{1,2}  · Olive M. Mbah¹ · Wendi Elkins¹ · Laura C. Pinheiro³ · Mary Anne Szymeczek¹ · Neda Padilla² · Jennifer S. Walker⁴ · Giselle Corbie-Smith⁵

Accepted: 11 May 2020

Abstract

Background Cancer is the leading cause of death among Hispanics/Latinos. Thus, understanding health-related quality of life (HRQOL) needs among this diverse racial/ethnic group is critical. Using Ferrell's multidimensional framework for measuring QOL, we synthesized evidence on HRQOL needs among Hispanic/Latino cancer survivors.

Methods We searched MEDLINE/PubMed, EMBASE, CINAHL, and PsycINFO, for English language articles published between 1995 and January 2020, reporting HRQOL among Hispanic/Latino cancer survivors in the USA.

Results Of the 648 articles reviewed, 176 met inclusion criteria, with 100 of these studies focusing exclusively on breast cancer patients and no studies examining end-of-life HRQOL issues. Compared with other racial/ethnic groups, Hispanics/Latinos reported lower HRQOL and a higher symptom burden across multiple HRQOL domains. Over 80% of studies examining racial/ethnic differences in psychological well-being ($n = 45$) reported worse outcomes among Hispanics/Latinos compared with other racial/ethnic groups. Hispanic/Latino cancer survivors were also more likely to report suboptimal physical well-being in 60% of studies assessing racial/ethnic differences ($n = 27$), and Hispanics/Latinos also reported lower social well-being relative to non-Hispanics/Latinos in 78% of studies reporting these outcomes ($n = 32$). In contrast, reports of spiritual well-being and spirituality-based coping were higher among Hispanics/Latinos cancer survivors in 50% of studies examining racial/ethnic differences ($n = 15$).

Discussion Findings from this review point to the need for more systematic and tailored interventions to address HRQOL needs among this growing cancer survivor population. Future HRQOL research on Hispanics/Latinos should evaluate variations in HRQOL needs across cancer types and Hispanic/Latino subgroups and assess HRQOL needs during metastatic and end-of-life disease phases.

Keywords Latino · Cancer survivor · Health-related quality of life · Supportive care

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s11136-020-02527-0>) contains supplementary material, which is available to authorized users.

Cleo A. Samuel
cleo_samuel@unc.edu

¹ Department of Health Policy and Management, Gillings School of Global Public Health, University of North Carolina, 1105F McGavran-Greenberg Hall, CB# 7411, Chapel Hill, NC 27599-7411, USA

² Lineberger Comprehensive Cancer Center, University of North Carolina, Chapel Hill, NC, USA

Background

Cancer is the leading cause of death among Hispanics/Latinos [1]. In the USA in 2018, there were approximately 150,000 incident cancer cases and 43,000 cancer-related deaths among Latinos [2]. For Hispanic/Latino men,

³ Department of General Internal Medicine, Weill Cornell Medical College, New York, NY, USA

⁴ University of North Carolina Health Sciences Library, University of North Carolina, Chapel Hill, NC, USA

⁵ Department of Social Medicine, School of Medicine, University of North Carolina, Chapel Hill, NC, USA

prostate, colorectal, and lung are the three most common cancers [2]; and among Hispanic/Latina women, the three most prevalent cancers are breast, thyroid, and uterine cancer [2]. Although Hispanics/Latinos have a 25% lower cancer incidence rate and 30% lower cancer mortality rate than non-Hispanic/Latino Whites; risk of infection-related cancers such as stomach, liver, and cervical cancer is nearly twice as high among Hispanics/Latinos [1]. Moreover, emerging evidence suggests that as immigration rates slow, and more Hispanics/Latinos are in the USA long-term, Hispanic/Latino incidence/mortality cancer rates will rise and become more similar to other racial/ethnic groups in the USA [3, 4]. These findings underscore the imperative need to better understand the cancer experience of this growing, heterogeneous US population.

Compared with non-Hispanic/Latino Whites, Hispanics/Latinos are more likely to report structural barriers to care including lower incomes and education, lack of health insurance, and no usual source of healthcare [5]. These socioeconomic differences may contribute to higher rates of late stage cancer diagnoses, and resultantly more aggressive treatment, among Hispanics/Latinos [2]. Aggressive cancer treatment modalities are known to pose significant symptom and quality of life burdens. Additionally, Hispanic/Latino cancer survivors face a range of psychosocial factors that may negatively affect their care quality and quality of life, including acculturation challenges, language barriers, and limited cancer self-efficacy [6–8]. In particular, language barriers have been linked to low patient satisfaction with their health care [9]. Research also suggests that Hispanics/Latinos without cancer are less likely to receive supportive care for mental health issues [10–13]. Thus, Hispanics/Latinos may be at increased risk for quality of life decrements following their cancer diagnoses and treatments.

Although previous systematic reviews summarized health-related quality of life (HRQOL) patterns among Latino cancer survivors, prior studies were largely focused on women with breast cancer, did not consider HRQOL experiences of survivors at different phases of the cancer care continuum, from the time of diagnosis to death, and/or failed to account for multiple domains of HRQOL [14–23]. Ferrell's Conceptual Framework on quality of life [24] has been used to study HRQOL among racially/ethnically diverse cancer survivors [24–26]. The framework considers four distinct HRQOL domains: 1) physical well-being (e.g., nausea, fatigue), social well-being (e.g., distress, sexual function), psychosocial well-being (e.g., anxiety, depression), and spiritual well-being (Fig. 1) [24]. The objective of this study was to synthesize evidence on HRQOL needs among Hispanic/Latino cancer survivors in the USA using Ferrell's HRQOL framework. This systematic review includes studies reporting HRQOL outcomes and needs among US-based Hispanic/Latino cancer survivors, of all

cancer types, from the time of diagnosis to death. This broad focus allows for a comprehensive assessment of the existing evidence on Hispanic/Latino HRQOL needs. Additionally, we present recommendations for advancing HRQOL research on Hispanic/Latino cancer survivors and improving supportive care services for this rapidly growing patient population.

Methods

Literature search strategy

A comprehensive literature search was conducted in MEDLINE/PubMed, EMBASE, CINAHL, and PsycINFO, to identify relevant articles published from 1995 (publication year of Ferrell's original article on *Measurement of QOL in Cancer Survivors* [24]) through January 2020. Search terms were optimized for each database and included Medical Subject Headings (MeSH) and related text/keyword searches when appropriate. The search focused on terms used to describe (a) HRQOL (e.g., financial burden, spirituality, depression, insomnia, sexual dysfunction); in (b) Hispanic/Latino (e.g., Hispanic, Latino, Puerto Rican, Cuban); (c) cancer survivors (e.g., cancer, carcinoma*, neoplasm*) AND (survivorship, survivor OR survivors). The full search strategy can be found in Supplemental Appendix Table 1.


Inclusion and exclusion criteria

Eligibility criteria were developed with respect to the population, outcomes of interest, study design, and timing (PICOTS), and publication type. Studies were eligible for inclusion if they presented results on HRQOL among Hispanic/Latino cancer survivors of any age. Consistent with current definitions of a “cancer survivor,” we employed a lifetime definition capturing time from cancer diagnosis until death from any cause [27, 28]. We used the US Census Bureau's definition of Hispanics/Latinos (individuals of Cuban, Mexican, Puerto Rican, South or Central American, or other Spanish culture or origin regardless of race [29]). Conference abstracts and non-peer-reviewed publications were excluded from this review, as were non-empirical studies (with the exception of literature reviews), non-English studies, non-US-based studies, and studies not separately reporting HRQOL for our target population (i.e., Hispanic/Latino cancer survivors).

Study selection

Two trained research assistants (RAs) used the eligibility criteria to independently screen titles and abstracts in duplicate for inclusion/exclusion. Studies with titles and abstracts that

Fig. 1 Ferrell's quality of life in cancer survivors



met the inclusion criteria or lacked adequate information to determine inclusion/exclusion underwent full-text review. During the full-text review, two RAs independently reviewed all full-text articles in duplicate for inclusion/exclusion. If both reviewers agreed that a study did not meet eligibility criteria, the study was excluded. A senior member of the review team resolved any conflicts. Covidence software was used for this review. The software was set to dual screening mode which required at least 2 reviewers to agree for article inclusion/exclusion at each review phase.

Data extraction and quality appraisal

Three independent reviewers abstracted data from the 176 eligible articles using a standardized abstraction form (Supplemental Appendix Table 2) that captured

study participant characteristics, conceptual frameworks employed, study design, HRQOL measures, and Latino-specific HRQOL findings. The three reviewers abstracted the first five articles together to ensure consistency. At least two reviewers were involved in each article abstraction – one person conducted the primary abstraction and the second person performed quality control. Reviewers also assessed the quality of each article using the 22-item Strengthening the Reporting of Observational Studies (STROBE) tool [30] for articles with quantitative methods, and the ten-item Critical Appraisal Skills Program (CASP) tool [31] for qualitative studies. Each study was assigned an appraisal tool ratio, calculated as the number of STROBE/CASP criteria met divided by the total number of tool criteria (Supplemental Appendix Table 2) [32]. No studies were excluded based on quality appraisal ratio scores.

Table 1 Summary of Included Studies

HRQOL domains	Study design	Examples of commonly explored HRQOL measures
Overall	Quantitative Cross-sectional: 25 studies [1–25] Quantitative Longitudinal: 7 studies [26–31] Qualitative: 1 study [32] Mixed-Methods: 0 studies Literature Review: 4 studies [33–36]	Overall: [2–8, 10, 11, 13–15, 17, 19–26, 32, 34–36]
Physical well-being	Quantitative Cross-sectional: 57 studies [2, 5, 6, 8, 9, 12–15, 17, 20, 21, 23, 24, 37–79] Quantitative Longitudinal: 21 studies [26, 28, 31, 80–97] Qualitative: 12 studies [98–108] Mixed-Methods: 7 studies [109–115] Literature Review: 8 studies [34–36, 116–120]	General Physical Health/ Functional Status: [5, 8, 14, 24, 28, 32, 38, 39, 41, 43, 46, 47, 50, 53, 54, 56, 59, 65, 68–70, 81, 83, 85, 90, 93, 94, 110–113] Lymphedema: [36, 51, 86, 97, 121] Pain: [51, 52, 69, 89, 101, 102, 106, 112] Fatigue: [28, 51, 64, 94, 103, 106]
Psychological well-being	Quantitative Cross-sectional: 74 studies [2–9, 12–18, 20, 21, 23–25, 37–44, 46, 48–65, 67–69, 71–76, 78, 122–138] Quantitative Longitudinal: 23 studies [26, 28, 31, 81–85, 87, 88, 90, 91, 93–96, 139–145] Qualitative: 27 studies [32, 98–108, 146–160] Mixed-Methods: 7 studies [109–112, 114, 115, 161] Literature Review: 12 studies [33–36, 116–120, 162–164]	General Mental Health/ Emotional Well-Being: [3, 5, 8, 24, 28, 39, 41, 43, 46, 50, 53, 54, 59, 65, 68, 69, 81, 83, 85, 90, 93, 110–113, 142] Coping: [3, 28, 102, 142, 155] Depression: [15, 28, 38, 64, 84, 94, 98, 102, 109, 122, 127, 139, 145] Stress: [4, 25, 28, 38, 98, 154]
Social well-being	Quantitative Cross-sectional: 55 studies [2, 3, 5, 7, 8, 12–16, 20–24, 37–42, 46, 48, 49, 53, 54, 56, 57, 59, 60, 62, 66, 67, 69, 73–76, 122–126, 129–132, 134, 135, 138, 165–169] Quantitative Longitudinal: 22 studies [28, 31, 81–84, 88, 90–94, 134, 140–142, 144, 145, 170–173] Qualitative: 25 studies [32, 98–106, 108, 147–152, 154, 156–160, 174, 175] Mixed-Methods: 5 studies [110–114] Literature Review: 10 studies [33–36, 116–118, 163, 164, 176]	Social Support: [15, 20, 22, 99, 101–104, 110, 114, 125, 135, 149, 150, 152, 154, 156–158] Family: [103, 106, 154] Finances: [4, 28, 32, 99, 101, 104, 106, 110, 132, 172, 173] Sexuality: [4, 28, 59, 84, 101, 103, 106]
Spiritual well-being	Quantitative Cross-sectional: 22 studies [5, 7, 8, 14, 16, 17, 20, 21, 24, 38–40, 47, 60, 64, 70, 123, 131, 135, 136, 177] Quantitative Longitudinal: 4 studies [28, 81, 94, 142] Qualitative: 24 studies [32, 98–106, 108, 147–150, 152–157, 160, 175, 178] Mixed-Methods: 3 studies [111–113] Literature Review: 6 studies [34, 35, 116, 118, 163, 164]	Spirituality/ Spiritual Well-Being: [14, 17, 24, 46, 81, 99, 100, 108, 112, 113, 135, 150, 156, 157, 178] Religion: [32, 102, 136, 150, 155, 175] Uncertainty/ Questioning God: [20, 21, 28] Prayer: [32, 64, 70, 94, 98, 106, 149]

Results


Literature search results

A total of 1,094 articles were identified through database searching, of which 648 were non-duplicates. A total of 377 articles were excluded during the initial title and abstract screening phase. Of the remaining 271 articles, 95 articles were excluded during full-text review, and 176 articles met all inclusion criteria (Fig. 2).

Description of included studies

The study designs of the 176 articles varied and included 90 (51.1%) cross-sectional quantitative, 35 (19.9%) longitudinal quantitative, 30 (17.0%) qualitative, eight (4.5%) mixed-methods, and 13 (7.4%) literature reviews (Table 1). One hundred studies focused exclusively on breast cancer, while the remainder included other (e.g., prostate) or multiple cancers. Time since cancer diagnosis for study participants ranged from 0–120 months

Fig. 2 Literature flow diagram



post-diagnosis. Most studies focused on adults. Forty studies employed a HRQOL conceptual framework and the majority of studies used validated HRQOL measurement instruments (e.g., Functional Assessment of Cancer Therapy–General [FACT-G]). All studies met at least 70% of the critical appraisal criteria, and 94 studies met > 70% of the appraisal criteria. Although literature reviews were included in this study, they were excluded from critical appraisal.

Findings by Health-Related Quality of Life Domain.

Results organized by Ferrell’s HRQOL domains (i.e., psychological, physical, social, and spiritual) are presented in Table 1 and Supplemental Appendix Table 2. We included an additional section labeled *Combined “Overall” HRQOL* that summarizes studies that included some combination of the four HRQOL domains [14]. In addition to reporting HRQOL patterns among Hispanic/Latino cancer survivors, several studies examined differences in HRQOL between Hispanic/Latino cancer survivors and survivors from other racial/ethnic groups and/or correlates of HRQOL. Such studies are summarized within each HRQOL domain sub-section

below. Where applicable, we summarized interventions to improve HRQOL among Latino cancer survivors.

Psychological well-being

Most included studies (143 out of 176 studies) focused on psychological well-being, with 38 studies reporting lower psychological well-being among Hispanics/Latinos compared to other racial/ethnic groups [6, 14–19, 22, 33–61]. Only six studies [62–67] reported similar mental health scores between Hispanics/Latinos cancer survivors and survivors from other racial/ethnic groups, and one study documented better mental health outcomes among less acculturated Hispanic/Latina breast cancer survivors compared to their White counterparts [68].

With a prevalence ranging from 32% [69] to 53% [70], depressive symptoms were the most commonly studied and reported HRQOL measure among Hispanic/Latino cancer survivors. Nine studies examined racial/ethnic differences in reports of depression, of which, six showed higher rates of depression among Hispanics/Latinos compared

to non-Hispanics/Latinos [15, 43, 52, 54, 57, 61], while the remaining three studies reported no ethnic differences [71–73]. For example, in a study examining post-treatment symptoms among breast cancer survivors, Hispanics/Latinas reported significantly higher rates of depression compared to non-Hispanic/Latino Whites and Blacks [43]. In contrast, a study examining HRQOL among cervical cancer survivors found no differences in rates of depression between Hispanics/Latinas and non-Hispanic/Latina survivors [73]. Several factors were associated with increased risk for depression among Hispanic/Latino cancer survivors, including lower education, poor physical/social functioning, less social (e.g., family, peer) and emotional support, higher stress, poor body image, poor coping, self-blame, and Spanish language preference [57, 70, 74, 75].

Other commonly studied and reported psychological well-being issues included emotional well-being concerns, distress or anxiety, and fear of recurrence [16–18, 34, 40, 50, 76–78]. Hispanic/Latino cancer survivors consistently reported worse emotional well-being compared to survivors of other racial/ethnic groups [14, 19, 37–39, 47–49, 51, 79, 80]. Of the 12 studies examining fear of cancer recurrence [16–18, 34, 40, 50, 58, 76–78, 81, 82], six reported higher levels of fear among Hispanics/Latinos compared to other racial/ethnic groups [16–18, 34, 40, 50, 78]. In a systematic review of studies on fear of recurrence among survivors of multiple cancers, Hispanic/Latino ethnicity was associated with higher levels of fear of cancer recurrence [18]. Additionally, two studies reported that Hispanics/Latinas with lower levels of acculturation reported the highest level of worry about recurrence compared to Hispanics/Latinas with high levels of acculturation, Black women, and White women [44, 45]. Other psychological HRQOL measures reported/studied among Hispanic/Latino cancer survivors included post-traumatic growth [35, 72, 83]; body image issues [36, 77, 78, 84–86]; stress [73, 87]; sadness [86]; worry about job loss or finances [16, 77, 88], worry about fertility, relationships, treatment effects [16, 22, 36, 51, 56, 84, 89]; neurocognitive performance [76, 90]; fear of social isolation [36]; coping [36, 88, 90–95]; and fatalism [96, 97].

Ten studies tested a variety of interventions to improve psychological well-being among Hispanic/Latino cancer survivors [20, 78, 98–105]. In-person and telephone-based education, as well as counseling and mindfulness-based stress reduction interventions were shown to improve anxiety, depressive symptoms, and psychological HRQOL among breast cancer survivors [78, 98, 101–103].

Physical well-being

A total of 105 studies focused on physical well-being among Hispanic/Latino cancer survivors. Twenty seven studies examined racial/ethnic differences in physical

well-being among Hispanic/Latino cancer survivors [6, 14, 37, 38, 41, 42, 47, 53–55, 62–68, 79, 106–114], with 16 of these studies documenting worse physical well-being among Hispanics/Latinos compared to survivors from other racial/ethnic groups [14, 38, 41, 42, 47, 53, 55, 65, 79, 106–110, 113, 114]. For example, two studies examining breast, colorectal, prostate and lung cancer survivors reported lower physical HRQOL scores among Hispanic/Latino survivors compared to other racial/ethnic groups [41, 42]. Acculturation was associated with physical HRQOL, where English proficient Hispanic/Latina breast cancer survivors were more likely than their low English proficient counterparts to report better self-rated physical health [6]. In contrast, eight studies reported no differences in physical HRQOL between Hispanic/Latino survivors and survivors from other racial/ethnic groups [54, 62–64, 66, 67, 111, 112]; while one study documented better physical well-being among less acculturated Hispanic/Latina breast cancer survivors relative to their non-Hispanic/Latina White counterparts [68].

Two studies evaluated racial/ethnic differences in physical functioning with conflicting findings. In a systematic review examining HRQOL among cancer and renal disease patients, Hispanic/Latino men with prostate cancer had significantly worse physical functioning than White men, and had more bowel-related issues than their White and Black counterparts [22]. Yet, in another study of breast, prostate, colorectal and gynecologic cancer patients, Latino cancer survivors reported higher physical functioning compared to Black and Asian cancer survivors [115].

Other measures of physical well-being reported among Hispanic/Latino cancer survivors included sexual functioning [16, 86, 116, 117], pain [14, 19, 43, 54, 77, 86, 91, 108, 118–122], fatigue [14, 19, 43, 54, 76, 78, 86, 104, 113, 119–121, 123–125], nausea [19, 86, 91, 96, 119, 121, 123], hair loss [19, 49, 86, 91, 123], lymphedema [14, 43], bowel problems [22, 123], and “other symptom bother” [49, 73, 91, 123, 126]. Studies examining racial/ethnic differences in sexual function consistently reported that Latino cancer survivors were more likely to experience adverse sexual function than other racial/ethnic groups [16, 17, 117]. Compared with other racial/ethnic groups, Hispanic/Latino and Black survivors reported worse physical toxicities from treatment [14, 22, 43, 49, 108, 117, 118]. Additionally, in a systematic review of HRQOL among Hispanic/Latina breast cancer patients, Hispanics/Latinas reported higher symptom burden for symptoms such as pain, fatigue and lymphedema [14].

Four studies examining the effect of telephone-based and multi-part interventions (including culturally sensitive health education and interpersonal counseling) on HRQOL among Hispanic/Latino cancer survivors reported significant improvements in physical well-being in intervention groups [20, 101, 104, 127].

Social well-being

One hundred and seventeen studies examined social well-being among Hispanic/Latino cancer survivors. Thirteen studies examined social well-being using a summary score measure (e.g., FACT-G) with nine reporting lower social well-being scores among Hispanics/Latinos compared to other racial/ethnic groups [14, 15, 17, 38, 39, 48, 53, 64, 109]. In a study examining HRQOL among a diverse group of cervical cancer survivors, Spanish-speaking Hispanics/Latinas survivors reported worse social well-being compared to White survivors [39]. In another cohort of multiethnic breast cancer survivors, Hispanics/Latinas reported the lowest scores for social/family well-being when compared to other groups [38]. Social support [16, 19, 36, 40, 50, 61, 63, 73, 76–78, 81, 82, 86, 88, 91–93, 95, 97, 102, 128–145] and financial hardship/job disruption [19, 33, 45, 50, 81, 82, 86, 102, 106, 121, 128, 130, 131, 140, 145–157] were the most commonly reported/studied social well-being measures. Hispanic/Latino cancer survivors reported social support as an important component of dealing with a cancer diagnosis [76–78, 81, 82, 86, 88, 92, 93, 95, 97, 102, 121, 131, 132, 134, 140, 145]. In a study among Hispanic/Latino adolescent and young adult cancer survivors, family communication around cancer and symptoms was reported as beneficial to transitioning these survivors from pediatric to adult-centered survivorship care [132]. While one study reported that Hispanic/Latino cancer survivors mainly sought support from family [21], another study noted that informal social support networks were more helpful to Hispanic/Latino survivors relative to their non-Hispanic/Latino White counterparts [16]. Although social support was strongly endorsed among Hispanic/Latino cancer survivors, several studies reported more adverse social HRQOL outcomes in Hispanic/Latino cancer survivors compared to other racial/ethnic groups, including smaller social support networks / less social support [19, 50, 138], more social avoidance [40], and lower scores on measures of interpersonal relationships [113, 139]. Hispanics/Latinos cancer survivors were also less likely to take part in support groups [135] and report changes in their marital status as a result of a cancer diagnosis [128]. In contrast, one study among cervical cancer patients reported no differences in social support between Hispanic/Latino and non-Hispanic/Latino survivors [73]. Moreover, in one study, while non-Hispanic/Latino Whites perceived more social support than Hispanics/Latinas (mean score of 145.9 vs. 139.0, $p=0.04$), Latinas reported slightly higher spousal and familial support [138].

Hispanic/Latino cancer survivors more commonly reported job disruptions and financial hardship after their cancer diagnosis [33, 50, 128, 147–152, 157]. For example, in a study examining correlates of paid work after breast cancer diagnosis, Hispanic/Latina breast cancer survivors were

more likely than Whites to stop working after a diagnosis [151]. However, another study examining factors associated with returning to work noted that there were no significant differences in job retention between Hispanic/Latino cancer survivors and Whites, although job retention was associated with acculturation [146]. Yet, another study found that Hispanic/Latino cancer survivors were more likely to return to work than White survivors [106]. Financial hardship or loss was also more prevalent among Hispanic/Latino cancer survivors compared to other racial/ethnic groups, with the exception of Blacks [86, 148, 149, 152, 157]. Other social HRQOL concerns reported among Hispanic/Latino cancer survivors included sexual health and sexuality (e.g., the negative effect of diagnosis on enjoying sex) [46, 71, 77, 84, 122, 142, 145].

Spiritual well-being

Fifty-nine studies examined spiritual well-being among Hispanic/Latino cancer survivors, and documented a range of religious/spiritual practices for coping with cancer, including: prayer [92, 111, 123, 158], faith [21, 36, 77, 84, 159, 160], finding meaning and peace [61, 143], religious/church attendance [158], and relationship with God [22, 47, 63, 64, 94, 111, 159–162]. Hispanic/Latino survivors commonly reported spirituality as a source of strength [78, 86, 93, 101] and comfort [47, 161]. The 15 studies [16, 22, 47, 61, 63, 64, 94, 111, 130, 143, 159–163] examining racial/ethnic differences in spiritual well-being generally showed that compared to other racial/ethnic groups, Hispanic/Latino and Black cancer survivors were more likely to use spirituality as a coping strategy [22, 61, 64, 94, 143, 160, 162]. In a study among patients with hematological diseases and brain tumors, compared with non-Hispanics/Latinos, Hispanic/Latino survivors reported higher spiritual well-being scores [162]. In contrast, a study examining HRQOL among survivors of childhood cancers showed no statistically significant difference in spiritual well-being among Hispanics/Latinos and non-Hispanic/Latino Whites [63]. Spirituality/religious coping was associated with other aspects of HRQOL among Hispanic/Latino cancer survivors [83, 158, 163], including lower anxiety, fatigue, and depression [158], better functional well-being [163], and less depression [61].

Combined “Overall” HRQOL

Thirty-seven studies reported on overall HRQOL domains among Hispanic/Latino cancer survivors, highlighting racial/ethnic differences between Hispanics/Latinos and other groups [8, 14, 15, 22, 39, 57, 65, 73, 75, 110, 162, 164–170]. In a multiethnic cohort of women with breast cancer, Hispanic/Latino ethnicity was associated with worse overall HRQOL compared to Black survivors [164]. Another

study of HRQOL among prostate cancer patients reported lower overall HRQOL among Hispanic/Latino and Black men relative to other racial/ethnic groups [22]. However, a study among cervical cancer survivors reported no differences in overall HRQOL between Hispanics/Latinas and non-Hispanics/Latinas [73]. In terms of correlates of HRQOL, greater acculturation, shame, and stigma were negatively associated with overall HRQOL, while more physical activity, inner peace, and better physical, social, psychological and functional well-being were positively associated with overall HRQOL among Hispanic/Latino cancer survivors [6, 57, 75, 110, 144, 163, 166–168].

Four studies documented efficacy of culturally sensitive education and counseling interventions, as well as patient navigation in improving overall HRQOL among Hispanic/Latino cancer patients [20, 78, 127, 171].

Discussion

In this systematic review, we synthesized evidence on HRQOL experiences and needs among Hispanic/Latino cancer survivors, from the time of cancer diagnosis until death. Evidence from our review suggests that HRQOL needs are high among Hispanic/Latino cancer survivors. In particular, Hispanics/Latinos reported worse psychological well-being (e.g., depression, anxiety, fear of recurrence) and suboptimal physical well-being (e.g., pain, nausea, sexual dysfunction) relative to their counterparts. In terms of social well-being, although Hispanic/Latino survivors commonly endorse preferences for family support and social network involvement during care, several studies reported lower social well-being and more job disruptions/financial hardship among Hispanics/Latinos relative to non-Hispanic/Latino Whites. In contrast, reports of spiritual well-being and spirituality-based coping were high among Hispanic/Latino cancer survivors.

Fewer studies evaluated HRQOL interventions; however, culturally responsive education and counseling were the most commonly tested interventions with demonstrated efficacy in improving HRQOL. However, given the extent of HRQOL needs in Hispanic/Latino cancer survivors, there is an urgent need for more systematic and tailored interventions to address the HRQOL needs of this growing and underrepresented segment of the cancer survivor population. In terms of potential intervention targets, previous research has documented racial/ethnic differences in clinicians' assessment of patients' symptoms, with clinicians being less likely to assess and accurately estimate HRQOL symptom burden in patients of color relative to Whites [172–175]. Uneven clinician screening of patients for HRQOL/symptom burden likely contributes to inequities in supportive care service access and delivery, including medication prescribing, and referrals to supportive care specialists (e.g., psychiatry,

pain specialist, physical therapy). Although distress screening has been an American College of Surgeons accreditation standard for cancer centers since 2015 [176], recent evidence suggests that the uptake and dissemination of distress screening has been limited in some settings, including National Cancer Institute-designated cancer centers [177]. Thus, efforts to improve HRQOL among Hispanic/Latino cancer survivors must include broader uptake of routine symptom and distress screening for HRQOL issues and supportive care needs, especially depression/anxiety, physical symptoms (e.g., pain, sexual dysfunction, bowel issues), job loss and financial hardship, social support, and spiritual coping. The National Comprehensive Cancer Network (NCCN) Distress Thermometer and Problem List is a helpful tool for quickly assessing both patient distress and a range of HRQOL concerns that closely align with the four HRQOL domains examined in this review [178]. The NCCN distress screening tool has been translated/adapted for Spanish-speakers which is important in this population due to heterogeneity in acculturation, English proficiency, and Spanish language preference. Furthermore, routine monitoring of symptom distress and HRQOL needs can be facilitated through the use of health information technology (e.g., electronic patient-reported outcome [ePRO] systems), where symptom distress and HRQOL needs can be addressed in real-time and tracked across visits. Prior studies have demonstrated the feasibility, acceptability, and efficacy of symptom monitoring through implementation of ePRO systems in routine oncology care [179], but feasibility/acceptability/efficacy testing needs to be expanded to more diverse cancer patient populations, including Hispanics/Latinos.

Moreover, addressing the HRQOL needs of Hispanic/Latino cancer survivors requires consideration of the unique cultural and linguistic needs of this patient population, as well as the current social and political context surrounding the lived experience of Hispanics/Latinos in America, where high uninsurance rates (19% of Hispanics/Latinos remain uninsured [180]) and rising anti-immigrant and xenophobic sentiments [181] continue to threaten the health and well-being of the Hispanic/Latino community. Specifically, clinicians should recognize that the socio-political context in which Hispanics/Latinos in America live may further exacerbate psychological distress and other HRQOL issues in many Hispanic/Latino survivors and hinder some from seeking and receiving critical follow-up care to address these needs. To advance equity in supportive care for Hispanic/Latino cancer survivors, it is important for cancer centers to provide culturally and linguistically appropriate supportive care services that include medical interpreters as a standard of care [182], support family/social network involvement in decision-making [182], account for patients' spiritual care needs in care planning [182], and acknowledge and address, where possible, the broader societal stressors (e.g.,

discrimination, financial hardship, job loss) that may affect Hispanic/Latino survivors' HRQOL and their engagement with the cancer care system.

While this systematic review resulted in over 150 articles meeting inclusion criteria, there are some limitations of the included studies that are worth noting. First, our study was limited to Latinos/Hispanics in the US context, where anti-immigration sentiments and health care coverage challenges remain key concerns for this subset of the cancer survivor population. Thus, our findings may not generalize to Latinos/Hispanic cancer survivors in other countries, especially countries where health care and immigration policies are more inclusive. Second, men were vastly underrepresented, with roughly a quarter of studies including men in their study sample and only two studies solely focused on men [82, 117]. Overrepresentation of women across studies potentially limits generalizability of our findings to men, which precludes efforts to understand and address gender-specific HRQOL needs among Hispanic/Latino men with cancer. Additionally, few studies presented results stratified by Hispanic/Latino subgroups (e.g., Mexican vs. Puerto Rican vs. Venezuelan), thereby overlooking possible heterogeneity in HRQOL needs among Hispanics/Latinos from different countries of origin [183]. Most studies also focused on patients with earlier stage diagnoses, but no study evaluated HRQOL needs of Hispanics/Latinos with metastatic disease or those near the end of life (EOL), when HRQOL needs are known to be greater. In terms of representation of HRQOL domains across studies, relatively few studies evaluated the social and spiritual aspects of HRQOL among Hispanics/Latinos. Finally, the majority of included studies involved a cross-sectional observational study design (90 out of 176 studies were cross-sectional; 35 out of 176 were longitudinal) which hinders identification of changes in HRQOL needs overtime, critical time frames for intervention, and effective strategies for enhancing HRQOL in this population. Future research on HRQOL among Latino cancer survivors should place greater emphasis on exploring the experiences of Latino cancer survivors in countries outside the USA, recruiting more men into studies, evaluating HRQOL needs during metastatic and EOL disease cases, stratifying data by Latino subgroup, including measures of social and spiritual HRQOL, and incorporating longitudinal and intervention study designs, where possible.

Conclusion

To our knowledge, this is the first systematic review to examine domain-specific HRQOL concerns and issues among Hispanic/Latino cancer survivors at different phases of the cancer care continuum, from cancer diagnosis to death. There is overwhelming evidence pointing to significant

HRQOL needs among Hispanic/Latino survivors, including depression/anxiety, decrements in physical function, job loss and financial hardship. Routine assessment of HRQOL needs in clinical practice, as well as culturally and linguistically responsive approaches to supportive cancer care hold much promise for enhancing HRQOL outcomes in this growing segment of the cancer survivor population.

Acknowledgements This work was supported by the National Cancer Institute (1 K01CA218473-01A1 to C.A.S). Olive M. Mbah is supported by the National Cancer Institute's National Research Service Award sponsored by the Lineberger Comprehensive Cancer Center at the University of North Carolina (T32 CA116339).

Author contributions Conceptualization: Cleo A. Samuel; Literature search and data analysis: Jennifer S. Walker, OMM, WE, MAS; Writing-original draft preparation: CAS, OMM, WE, LP, JSW, MAS; Writing- review and editing: CAS, OMM, WE, LP, GCS; Project Administration: NRP.

Compliance with ethical standards

Conflict of interest Dr. Cleo A. Samuel reports research funding from National Cancer Institute and Pfizer for work unrelated to this review. All other authors do not have any conflicts of interest to disclose.

References

1. Miller, K. D., Goding Sauer, A., Ortiz, A. P., Fedewa, S. A., Pinheiro, P. S., Tortolero-Luna, G., et al. (2018). Cancer Statistics for Hispanics/Latinos 2018. *CA: A Cancer Journal of Clinicians*, 68(6), 425–445. <https://doi.org/10.3322/caac.21494>.
2. American Cancer Society. (2018). *Cancer facts & figures for Hispanics/Latinos 2018–2020*. Atlanta: American Cancer Society, Inc.
3. Pinheiro, P. S., Sherman, R. L., Trapido, E. J., Fleming, L. E., Huang, Y., Gomez-Marin, O., et al. (2009). Cancer incidence in first generation US Hispanics: Cubans, Mexicans, Puerto Ricans, and new Latinos. *Cancer Epidemiology Biomarkers Prevention*, 18(8), 2162–2169. <https://doi.org/10.1158/1055-9965.epi-09-0329>.
4. Pinheiro, P. S., Callahan, K. E., Stern, M. C., & de Vries, E. (2018). Migration from Mexico to the United States: A high-speed cancer transition. *International Journal of Cancer*, 142(3), 477–488. <https://doi.org/10.1002/ijc.31068>.
5. Vélez, L., Chalela, P., & Ramirez, A. (2008). Hispanic/Latino health and disease: An overview. *Health Promotion in Multicultural Populations: A Handbook for Practitioners and Students*.
6. Garcia-Jimenez, M., Santoyo-Olsson, J., Ortiz, C., Lahiff, M., Sokal-Gutierrez, K., & Napoles, A. M. (2014). Acculturation, inner peace, cancer self-efficacy, and self-rated health among Latina breast cancer survivors. *Journal of Health Care for the Poor and Underserved*, 25(4), 1586–1602. <https://doi.org/10.1353/hpu.2014.0158>.
7. Kagawa-Singer, M. (1995). Socioeconomic and cultural influences on cancer care of women. *Seminars in Oncology Nursing*, 11(2), 109–119.
8. Medeiros, E. A., Castaneda, S. F., Gonzalez, P., Rodriguez, B., Buelna, C., West, D., et al. (2015). Health-related quality of life among cancer survivors attending support Groups. *Journal of*

- Cancer Education*, 30(3), 421–427. <https://doi.org/10.1007/s13187-014-0697-8>.
9. Baker, D. W., Hayes, R., & Fortier, J. P. (1998). Interpreter use and satisfaction with interpersonal aspects of care for Spanish-speaking patients. *Medical Care*, 36(10), 1461–1470.
10. Akincigil, A., Olsson, M., Siegel, M., Zurlo, K. A., Walkup, J. T., & Crystal, S. (2012). Racial and ethnic disparities in depression care in community-dwelling elderly in the United States. *American Journal of Public Health*, 102(2), 319–328. <https://doi.org/10.2105/ajph.2011.300349>.
11. Alegria, M., Canino, G., Rios, R., Vera, M., Calderon, J., Rusch, D., et al. (2002). Inequalities in use of specialty mental health services among Latinos, African Americans, and non-Latino whites. *Psychiatric Services (Washington, DC)*, 53(12), 1547–1555. <https://doi.org/10.1176/appi.ps.53.12.1547>.
12. Alegria, M., Chatterji, P., Wells, K., Cao, Z., Chen, C. N., Takeuchi, D., et al. (2008). Disparity in depression treatment among racial and ethnic minority populations in the United States. *Psychiatric Services (Washington, DC)*, 59(11), 1264–1272. <https://doi.org/10.1176/appi.ps.59.11.1264>.
13. Lagomasino, I. T., Stockdale, S. E., & Miranda, J. (2011). Racial-ethnic composition of provider practices and disparities in treatment of depression and anxiety, 2003–2007. *Psychiatric Services (Washington, DC)*, 62(9), 1019–1025. <https://doi.org/10.1176/appi.ps.62.9.1019>.
14. Yanez, B., Thompson, E. H., & Stanton, A. L. (2011). Quality of life among Latina breast cancer patients: A systematic review of the literature. *Journal of Cancer Survivorship*, 5(2), 191–207. <https://doi.org/10.1007/s11764-011-0171-0>.
15. Lockett, T., Goldstein, D., Butow, P. N., Gebiski, V., Aldridge, L. J., McGrane, J., et al. (2011). Psychological morbidity and quality of life of ethnic minority patients with cancer: A systematic review and meta-analysis. *The Lancet Oncology*, 12(13), 1240–1248. [https://doi.org/10.1016/s1470-2045\(11\)70212-1](https://doi.org/10.1016/s1470-2045(11)70212-1).
16. Aziz, N. M., & Rowland, J. H. (2002). Cancer survivorship research among ethnic minority and medically underserved groups. *Oncology Nursing Forum*, 29(5), 789–801. <https://doi.org/10.1188/02.onf.789-801>.
17. Blinder, V. S., & Griggs, J. J. (2013). Health disparities and the cancer survivor. *Seminars in Oncology*, 40(6), 796–803. <https://doi.org/10.1053/j.seminoncol.2013.09.003>.
18. Koch, L., Jansen, L., Brenner, H., & Arndt, V. (2013). Fear of recurrence and disease progression in long-term (>= 5 years) cancer survivors: A systematic review of quantitative studies. *Psychooncology*, 22(1), 1–11. <https://doi.org/10.1002/pon.3022>.
19. Lopez-Class, M., Gomez-Duarte, J., Graves, K., & Ashing-Giwa, K. (2012). A contextual approach to understanding breast cancer survivorship among Latinas. *Psychooncology*, 21(2), 115–124. <https://doi.org/10.1002/pon.1998>.
20. McNulty, J., Kim, W., Thurston, T., Kim, J., & Larkey, L. (2016). Interventions to improve quality of life, well-being, and care in latino cancer survivors: A systematic literature review. *Oncology Nursing Forum*, 43(3), 374–384. <https://doi.org/10.1188/16.onf.374-384>.
21. Pfaendler, K. S., Wenzel, L., Mechanic, M. B., & Penner, K. R. (2015). Cervical cancer survivorship: Long-term quality of life and social support. *Clinical Therapeutics*, 37(1), 39–48. <https://doi.org/10.1016/j.clinthera.2014.11.013>.
22. Wang, J., Sallfrank, M. F., Hong, S. H., Liu, C., & Gourley, D. R. (2007). Patterns of reporting health-related quality of life across racial and ethnic groups. *Expert Review of Pharmacoeconomics and Outcomes Research*, 7(2), 177–186. <https://doi.org/10.1586/14737167.7.2.177>.
23. Yoo, G. J., Levine, E. G., & Pasick, R. (2014). Breast cancer and coping among women of color: A systematic review of the literature. *Supportive Care in Cancer*, 22(3), 811–824. <https://doi.org/10.1007/s00520-013-2057-3>.
24. Ferrell, B. R., Dow, K. H., & Grant, M. (1995). Measurement of the quality of life in cancer survivors. *Quality of Life Research*, 4(6), 523–531.
25. Burhansstipanov, L., Dignan, M., Jones, K. L., Krebs, L. U., Marchionda, P., & Kaur, J. S. (2012). Comparison of quality of life between Native and non-Native cancer survivors: Native and non-Native cancer survivors' QOL. *Journal of Cancer Education*, 27(1 Suppl), S106–S113. <https://doi.org/10.1007/s13187-012-0318-3>.
26. Samuel, C. A., Pinheiro, L. C., Reeder-Hayes, K. E., Walker, J. S., Corbie-Smith, G., Fashaw, S. A., et al. (2016). To be young, Black, and living with breast cancer: A systematic review of health-related quality of life in young Black breast cancer survivors. *Breast Cancer Research and Treatment*, 160(1), 1–15. <https://doi.org/10.1007/s10549-016-3963-0>.
27. American Cancer Society. (2014). *Life after cancer: Survivorship by the numbers*. Atlanta: American Cancer Society, Inc.
28. Cho, J., Jung, S. Y., Lee, J. E., Shim, E. J., Kim, N. H., Kim, Z., et al. (2014). A review of breast cancer survivorship issues from survivors' perspectives. *Journal of Breast Cancer*, 17(3), 189–199. <https://doi.org/10.4048/jbc.2014.17.3.189>.
29. U.S. Office of Management and Budget; United States Census Bureau. Hispanic Origin. 2019. Retrieved July 2, 2019.
30. von Elm, E., Altman, D. G., Egger, M., Pocock, S. J., Gøtzsche, P. C., & Vandenbroucke, J. P. (2008). The Strengthening of Reporting of Observational Studies in Epidemiology (STROBE) statement: Guidelines for reporting observational studies. *Journal of Clinical Epidemiology*, 61(4), 344–349. <https://doi.org/10.1016/j.jclinepi.2007.11.008>.
31. Critical Appraisal Skills Program. CASP Qualitative Checklist. (2014).
32. Mollica, M., Nemeth, L., Newman, S. D., & Mueller, M. (2015). Quality of life in African American Breast Cancer Survivors: An integrative literature review. *Cancer Nursing*, 38(3), 194–204. <https://doi.org/10.1097/ncc.0000000000000160>.
33. Bowen, D. J., Alfano, C. M., McGregor, B. A., Kuniyuki, A., Bernstein, L., Meeske, K., et al. (2007). Possible socioeconomic and ethnic disparities in quality of life in a cohort of breast cancer survivors. *Breast Cancer Research and Treatment*, 106(1), 85–95. <https://doi.org/10.1007/s10549-006-9479-2>.
34. Blinder, V. S., Murphy, M. M., Vahdat, L. T., Gold, H. T., de Melo-Martin, I., Hayes, M. K., et al. (2012). Employment after a breast cancer diagnosis: A qualitative study of ethnically diverse urban women. *Journal of Community Health*, 37(4), 763–772. <https://doi.org/10.1007/s10900-011-9509-9>.
35. Arpawong, T. E., Oland, A., Milam, J. E., Ruccione, K., & Meeske, K. A. (2013). Post-traumatic growth among an ethnically diverse sample of adolescent and young adult cancer survivors. *Psychooncology*, 22(10), 2235–2244. <https://doi.org/10.1002/pon.3286>.
36. Ashing-Giwa, K. T., Kagawa-Singer, M., Padilla, G. V., Tejero, J. S., Hsiao, E., Chhabra, R., et al. (2004). The impact of cervical cancer and dysplasia: A qualitative, multiethnic study. *Psychooncology*, 13(10), 709–728. <https://doi.org/10.1002/pon.785>.
37. Ashing-Giwa, K. T., & Lim, J. W. (2011). Examining emotional outcomes among a multiethnic cohort of breast cancer survivors. *Oncology Nursing Forum*, 38(3), 279–288. <https://doi.org/10.1188/11.onf.279-288>.
38. Ashing-Giwa, K. T., Tejero, J. S., Kim, J., Padilla, G. V., & Hellemann, G. (2007). Examining predictive models of HRQOL in a population-based, multiethnic sample of women with breast carcinoma. *Quality of Life Research*, 16(3), 413–428. <https://doi.org/10.1007/s11136-006-9138-4>.

39. Ashing-Giwa, K. T., Tejero, J. S., Kim, J., Padilla, G. V., Kagawa-Singer, M., Tucker, M. B., et al. (2009). Cervical cancer survivorship in a population based sample. *Gynecologic Oncology*, 112(2), 358–364. <https://doi.org/10.1016/j.ygyno.2008.11.002>.
40. Carver, C. S., Smith, R. G., Petronis, V. M., & Antoni, M. H. (2006). Quality of life among long-term survivors of breast cancer: Different types of antecedents predict different classes of outcomes. *Psychooncology*, 15(9), 749–758. <https://doi.org/10.1002/pon.1006>.
41. Clauser, S. B., Arora, N. K., Bellizzi, K. M., Haffer, S. C., Topor, M., & Hays, R. D. (2008). Disparities in HRQOL of cancer survivors and non-cancer managed care enrollees. *Health Care and Finance Review*, 29(4), 23–40.
42. Connor, A. E., Baumgartner, R. N., Pinkston, C. M., Boone, S. D., & Baumgartner, K. B. (2016). Obesity, ethnicity, and quality of life among breast cancer survivors and women without breast cancer: The long-term quality of life follow-up study. *Cancer Causes and Control*, 27(1), 115–124. <https://doi.org/10.1007/s10552-015-0688-z>.
43. Eversley, R., Estrin, D., Dibble, S., Wardlaw, L., Pedrosa, M., & Favila-Penney, W. (2005). Post-treatment symptoms among ethnic minority breast cancer survivors. *Oncology Nursing Forum*, 32(2), 250–256. <https://doi.org/10.1188/05.onf.250-256>.
44. Janz, N. K., Hawley, S. T., Mujahid, M. S., Griggs, J. J., Alderman, A., Hamilton, A. S., et al. (2011). Correlates of worry about recurrence in a multiethnic population-based sample of women with breast cancer. *Cancer*, 117(9), 1827–1836. <https://doi.org/10.1002/cncr.25740>.
45. Janz, N. K., Li, Y., Beesley, L. J., Wallner, L. P., Hamilton, A. S., Morrison, R. A., et al. (2016). Worry about recurrence in a multi-ethnic population of breast cancer survivors and their partners. *Supportive Care in Cancer*, 24(11), 4669–4678. <https://doi.org/10.1007/s00520-016-3314-z>.
46. Knopf, M. T., Ferrucci, L. M., Cartmel, B., Jones, B. A., Stevens, D., Smith, M., et al. (2012). Needs assessment of cancer survivors in Connecticut. *Journal of Cancer Survivorship*, 6(1), 1–10. <https://doi.org/10.1007/s11764-011-0198-2>.
47. Lim, J. W., Gonzalez, P., Wang-Letzky, M. F., & Ashing-Giwa, K. T. (2009). Understanding the cultural health belief model influencing health behaviors and health-related quality of life between Latina and Asian-American breast cancer survivors. *Supportive Care in Cancer*, 17(9), 1137–1147. <https://doi.org/10.1007/s00520-008-0547-5>.
48. Meeske, K. A., Patel, S. K., Palmer, S. N., Nelson, M. B., & Parow, A. M. (2007). Factors associated with health-related quality of life in pediatric cancer survivors. *Pediatric Blood & Cancer*, 49(3), 298–305. <https://doi.org/10.1002/pbc.20923>.
49. Munoz, A. R., Kaiser, K., Yanez, B., Victorson, D., Garcia, S. F., Snyder, M. A., et al. (2016). Cancer experiences and health-related quality of life among racial and ethnic minority survivors of young adult cancer: A mixed methods study. *Supportive Care in Cancer*, 24(12), 4861–4870. <https://doi.org/10.1007/s00520-016-3340-x>.
50. Pandya, D. M., Patel, S., Ketchum, N. S., Pollock, B. H., & Padmanabhan, S. (2011). A comparison of races and leukemia subtypes among patients in different cancer survivorship phases. *Clin Lymphoma Myeloma Leuk*, 11(Suppl 1), S114–S118. <https://doi.org/10.1016/j.clml.2011.05.036>.
51. Pinheiro, L. C., Wheeler, S. B., Chen, R. C., Mayer, D. K., Lyons, J. C., & Reeve, B. B. (2015). The effects of cancer and racial disparities in health-related quality of life among older Americans: A case-control, population-based study. *Cancer*, 121(8), 1312–1320. <https://doi.org/10.1002/cncr.29205>.
52. Wu, C., Ashing, K. T., Jones, V. C., & Barcelo, L. (2017). The association of neighborhood context with health outcomes among ethnic minority breast cancer survivors. *Journal of Behavioral Medicine*. <https://doi.org/10.1007/s10865-017-9875-6>.
53. Ashing, K. T., George, M., & Jones, V. (2018). Health-related quality of life and care satisfaction outcomes: Informing psychosocial oncology care among Latina and African-American young breast cancer survivors. *Psychooncology*, 27(4), 1213–1220. <https://doi.org/10.1002/pon.4650>.
54. Bevilacqua, L. A., Dulak, D., Schofield, E., Starr, T. D., Nelson, C. J., Roth, A. J., et al. (2018). Prevalence and predictors of depression, pain, and fatigue in older- versus younger-adult cancer survivors. *Psychooncology*, 27(3), 900–907. <https://doi.org/10.1002/pon.4605>.
55. Olagunju, T. O., Liu, Y., Liang, L. J., Stomber, J. M., Griggs, J. J., Ganz, P. A., et al. (2018). Disparities in the survivorship experience among Latina survivors of breast cancer. *Cancer*, 124(11), 2373–2380. <https://doi.org/10.1002/cncr.31342>.
56. Papaleontiou, M., Reyes-Gastelum, D., Gay, B. L., Ward, K. C., Hamilton, A. S., Hawley, S. T., et al. (2019). Worry in thyroid cancer survivors with a favorable prognosis. *Thyroid*, 29(8), 1080–1088.
57. Ritt-Olson, A., Miller, K., Baezconde-Garbanati, L., Freyer, D., Ramirez, C., Hamilton, A., et al. (2018). Depressive symptoms and quality of life among adolescent and young adult cancer survivors: Impact of gender and Latino culture. *Journal of Adolescent and Young Adult Oncology*, 7(3), 384–388.
58. Séguin Leclair, C., Lebel, S., & Westmaas, J. L. (2019). The relationship between fear of cancer recurrence and health behaviors: A nationwide longitudinal study of cancer survivors. *Health Psychology*, 38(7), 596.
59. StGeorge, S. M., Noriega Esquivas, B., Agosto, Y., Kobayashi, M., Leite, R., Vanegas, D., et al. (2020). Development of a multigenerational digital lifestyle intervention for women cancer survivors and their families. *Psycho-Oncology*, 29(1), 182–194.
60. Young, K., Shliakhtsitsava, K., Natarajan, L., Myers, E., Dietz, A. C., Gorman, J. R., et al. (2019). Fertility counseling before cancer treatment and subsequent reproductive concerns among female adolescent and young adult cancer survivors. *Cancer*, 125(6), 980–989. <https://doi.org/10.1002/cncr.31862>.
61. Yu, Q., Medeiros, K. L., Wu, X., & Jensen, R. E. (2018). Non-linear predictive models for multiple mediation analysis: With an application to explore ethnic disparities in anxiety and depression among cancer survivors. *Psychometrika*, 83(4), 991–1006. <https://doi.org/10.1007/s11336-018-9612-2>.
62. Reyes, M. E., Ye, Y., Zhou, Y., Liang, A., Kopetz, S., Rodriguez, M. A., et al. (2017). Predictors of health-related quality of life and association with survival may identify colorectal cancer patients at high risk of poor prognosis. *Quality of Life Research*, 26(2), 319–330. <https://doi.org/10.1007/s11136-016-1381-8>.
63. Casillas, J. N., Zebrack, B. J., & Zeltzer, L. K. (2006). Health-related quality of life for Latino survivors of childhood cancer. *Journal of Psychosocial Oncology*, 24(3), 125–145. https://doi.org/10.1300/J077v24n03_06.
64. Bloom, J. R., Stewart, S. L., Napoles, A. M., Hwang, E. S., Livaudais, J. C., Karliner, L., et al. (2013). Quality of life of Latina and Euro-American women with ductal carcinoma in situ. *Psychooncology*, 22(5), 1008–1016. <https://doi.org/10.1002/pon.3098>.
65. Castellino, S. M., Casillas, J., Hudson, M. M., Mertens, A. C., Whitton, J., Brooks, S. L., et al. (2005). Minority adult survivors of childhood cancer: A comparison of long-term outcomes, health care utilization, and health-related behaviors from the childhood cancer survivor study. *Journal of Clinical Oncology*, 23(27), 6499–6507. <https://doi.org/10.1200/jco.2005.11.098>.
66. Pakiz, B., Ganz, P. A., Sedjo, R. L., Flatt, S. W., Demark-Wahnefried, W., Liu, J., et al. (2016). Correlates of quality of life in overweight or obese breast cancer survivors at enrollment into a

- weight loss trial. *Psychooncology*, 25(2), 142–149. <https://doi.org/10.1002/pon.3820>.
67. Goldfarb, M., & Casillas, J. (2016). Thyroid cancer-specific quality of life and health-related quality of life in young adult thyroid cancer survivors. *Thyroid*, 26(7), 923–932. <https://doi.org/10.1089/thy.2015.0589>.
68. Maly, R. C., Liu, Y., Liang, L. J., & Ganz, P. A. (2014). Quality of life over 5 years after a breast cancer diagnosis among low-income women: Effects of race/ethnicity and patient-physician communication. *Cancer*, 121(6), 916–926. <https://doi.org/10.1002/cncr.29150>.
69. Holden, A. E., Ramirez, A. G., & Gallion, K. (2014). Depressive symptoms in Latina breast cancer survivors: A barrier to cancer screening. *Health Psychology*, 33(3), 242–248. <https://doi.org/10.1037/a0032685>.
70. Ashing-Giwa, K., Rosales, M., Lai, L., & Weitzel, J. (2013). Depressive symptomatology among Latina breast cancer survivors. *Psychooncology*, 22(4), 845–853. <https://doi.org/10.1002/pon.3084>.
71. Christie, K. M., Meyerowitz, B. E., & Maly, R. C. (2010). Depression and sexual adjustment following breast cancer in low-income Hispanic and non-Hispanic White women. *Psychooncology*, 19(10), 1069–1077. <https://doi.org/10.1002/pon.1661>.
72. Foley, K. L., Farmer, D. F., Petronis, V. M., Smith, R. G., McGraw, S., Smith, K., et al. (2006). A qualitative exploration of the cancer experience among long-term survivors: Comparisons by cancer type, ethnicity, gender, and age. *Psychooncology*, 15(3), 248–258. <https://doi.org/10.1002/pon.942>.
73. Osann, K., Hsieh, S., Nelson, E. L., Monk, B. J., Chase, D., Cella, D., et al. (2014). Factors associated with poor quality of life among cervical cancer survivors: Implications for clinical care and clinical trials. *Gynecologic Oncology*, 135(2), 266–272. <https://doi.org/10.1016/j.ygyno.2014.08.036>.
74. Aguado Loi, C. X., Baldwin, J. A., McDermott, R. J., McMillan, S., Martinez Tyson, D., Yampolskaya, S., et al. (2013). Risk factors associated with increased depressive symptoms among Latinas diagnosed with breast cancer within 5 years of survivorship. *Psychooncology*, 22(12), 2779–2788. <https://doi.org/10.1002/pon.3357>.
75. Tobin, J., Miller, K. A., Baezconde-Garbanati, L., Unger, J. B., Hamilton, A. S., Milam, J. E., et al. (2018). Acculturation, mental health, and quality of life among hispanic childhood cancer survivors: A latent class analysis. *Ethnicity & Disease*, 28(1), 55–60. <https://doi.org/10.18865/ed.28.1.55>.
76. Ceballos, R. M., Molina, Y., Malen, R. C., Ibarra, G., Escareno, M., & Marchello, N. (2015). Design, development, and feasibility of a spanish-language cancer survivor support group. *Supportive Care in Cancer*, 23(7), 2145–2155. <https://doi.org/10.1007/s00520-014-2549-9>.
77. Ashing-Giwa, K. T., Padilla, G. V., Bohorquez, D. E., Tejero, J. S., Garcia, M., & Meyers, E. A. (2006). Survivorship: A qualitative investigation of Latinas diagnosed with cervical cancer. *Journal of Psychosocial Oncology*, 24(4), 53–88. https://doi.org/10.1300/J077v24n04_04.
78. Juarez, G., Hurria, A., Uman, G., & Ferrell, B. (2013). Impact of a bilingual education intervention on the quality of life of Latina breast cancer survivors. *Oncology Nursing Forum*, 40(1), E50–60. <https://doi.org/10.1188/13.onf.e50-e60>.
79. Giedzinska, A. S., Meyerowitz, B. E., Ganz, P. A., & Rowland, J. H. (2004). Health-related quality of life in a multiethnic sample of breast cancer survivors. *Annals of Behavioral Medicine*, 28(1), 39–51. https://doi.org/10.1207/s15324796abm2801_6.
80. Ashing-Giwa, K. T., & Lim, J. W. (2009). Examining the impact of socioeconomic status and socioecologic stress on physical and mental health quality of life among breast cancer survivors. *Oncology Nursing Forum*, 36(1), 79–88. <https://doi.org/10.1188/09.onf.79-88>.
81. Borrayo, E. A., Scott, K. L., Drennen, A., Bendriss, T. M., Kilbourn, K. M., & Valverde, P. (2020). Treatment challenges and support needs of underserved Hispanic patients diagnosed with lung cancer and head-and-neck cancer. *Journal of Psychosocial Oncology*, 2020, 1–14.
82. Sommariva, S., Vázquez-Otero, C., Medina-Ramirez, P., Aguado Loi, C., Fross, M., Dias, E., et al. (2019). Hispanic male cancer survivors' coping strategies. *Hispanic Journal of Behavioral Sciences*, 41(2), 267–284. <https://doi.org/10.1177/0739986319840658>.
83. Tobin, J., Allem, J. P., Slaughter, R., Unger, J. B., Hamilton, A. S., & Milam, J. E. (2017). Posttraumatic growth among childhood cancer survivors: Associations with ethnicity, acculturation, and religious service attendance. *Journal of Psychological Oncology*, 2017, 1–14. <https://doi.org/10.1080/07347332.2017.1365799>.
84. Ashing-Giwa, K. T., Padilla, G. V., Bohorquez, D. E., Tejero, J. S., & Garcia, M. (2006). Understanding the breast cancer experience of Latina women. *Journal of Psychosocial Oncology*, 24(3), 19–52. https://doi.org/10.1300/J077v24n03_02.
85. Buki, L. P., Reich, M., & Lehardy, E. N. (2016). "Our organs have a purpose": body image acceptance in Latina breast cancer survivors. *Psychooncology*, 25(11), 1337–1342. <https://doi.org/10.1002/pon.4270>.
86. Fatone, A. M., Moadel, A. B., Foley, F. W., Fleming, M., & Jandorf, L. (2007). Urban voices: the quality-of-life experience among women of color with breast cancer. *Palliat Support Care*, 5(2), 115–125.
87. Martinez Tyson, D. D., Jacobsen, P., & Meade, C. D. (2016). Understanding the stress management needs and preferences of Latinas undergoing chemotherapy. *Journal of Cancer Education*, 31(4), 633–639. <https://doi.org/10.1007/s13187-015-0844-x>.
88. Phillips, F., & Jones, B. L. (2014). Understanding the lived experience of Latino adolescent and young adult survivors of childhood cancer. *Journal of Cancer Survivorship*, 8(1), 39–48. <https://doi.org/10.1007/s11764-013-0310-x>.
89. Costas-Muniz, R., Hunter-Hernandez, M., Garduno-Ortega, O., Morales-Cruz, J., & Gany, F. (2017). Ethnic differences in psychosocial service use among non-Latina white and Latina breast cancer survivors. *Journal of Psychosocial Oncology*, 35(4), 424–437. <https://doi.org/10.1080/07347332.2017.1310167>.
90. Ayala-Feliciano, M., Pons-Valerio, J. J., Pons-Madera, J., & Acevedo, S. F. (2011). The relationship between visuospatial memory and coping strategies in breast cancer survivors. *Breast Cancer (Auckland)*, 5, 117–130. <https://doi.org/10.4137/bcbr.s6957>.
91. Banas, J. R., Victorson, D., Gutierrez, S., Cordero, E., Guitleman, J., & Haas, N. (2017). Developing a peer-to-peer mhealth application to connect Hispanic cancer patients. *Journal of Cancer Education*, 32(1), 158–165. <https://doi.org/10.1007/s13187-016-1066-6>.
92. Campesino, M. (2009). Exploring perceptions of cancer care delivery among older Mexican American adults. *Oncology Nursing Forum*, 36(4), 413–420. <https://doi.org/10.1188/09.onf.413-420>.
93. Carrion, I. V., Nedjat-Haiem, F., Macip-Billbe, M., & Black, R. (2017). "I Told Myself to Stay Positive" perceptions of coping among Latinos with a cancer diagnosis living in the United States. *American Journal of Hospital Palliative Care*, 34(3), 233–240. <https://doi.org/10.1177/1049909115625955>.
94. Gonzalez, P., Nunez, A., Wang-Letzkus, M., Lim, J. W., Flores, K. F., & Napoles, A. M. (2016). Coping with breast cancer: Reflections from Chinese American, Korean American, and

- Mexican American women. *Health Psychology*, 35(1), 19–28. <https://doi.org/10.1037/hea0000263>.
95. Jones, B. L., Volker, D. L., Vinajeras, Y., Butros, L., Fitchpatrick, C., & Rossetto, K. (2010). The meaning of surviving cancer for Latino adolescents and emerging young adults. *Cancer Nursing*, 33(1), 74–81. <https://doi.org/10.1097/NCC.0b013e3181b4ab8f>.
 96. Buki, L. P., Garces, D. M., Hiestrosa, M. C., Kogan, L., Carrillo, I. Y., & French, B. (2008). Latina breast cancer survivors' lived experiences: Diagnosis, treatment, and beyond. *Culture Diversity Ethnic Minor Psychology*, 14(2), 163–167. <https://doi.org/10.1037/1099-9809.14.2.163>.
 97. Livaudais, J. C., Thompson, B., Godina, R., Islas, I., Ibarra, G., & Coronado, G. D. (2010). A qualitative investigation of cancer survivorship experiences among rural Hispanics. *Journal of Psychosocial Oncology*, 28(4), 361–380. <https://doi.org/10.1080/07347332.2010.488146>.
 98. Ashing, K., & Rosales, M. (2014). A telephonic-based trial to reduce depressive symptoms among Latina breast cancer survivors. *Psychooncology*, 23(5), 507–515. <https://doi.org/10.1002/pon.3441>.
 99. Hughes, D. C., Leung, P., & Naus, M. J. (2008). Using single-system analyses to assess the effectiveness of an exercise intervention on quality of life for Hispanic breast cancer survivors: A pilot study. *Social Work in Health Care*, 47(1), 73–91. <https://doi.org/10.1080/00981380801970871>.
 100. Molina, Y., Thompson, B., Espinoza, N., & Ceballos, R. (2013). Breast cancer interventions serving US-based Latinas: current approaches and directions. *Womens Health (London)*, 9(4), 335–348. <https://doi.org/10.2217/whe.13.30>.
 101. Badger, T. A., Segrin, C., Hepworth, J. T., Pasvogel, A., Weihs, K., & Lopez, A. M. (2013). Telephone-delivered health education and interpersonal counseling improve quality of life for Latinas with breast cancer and their supportive partners. *Psychooncology*, 22(5), 1035–1042. <https://doi.org/10.1002/pon.3101>.
 102. Badger, T. A., Segrin, C., Sikorskii, A., Pasvogel, A., Weihs, K., Lopez, A. M., et al. (2020). Randomized controlled trial of supportive care interventions to manage psychological distress and symptoms in Latinas with breast cancer and their informal caregivers. *Psychology & Health*, 35(1), 87–106.
 103. Elimimian, E., Elson, L., Bilani, N., Farrag, S. E., Dwivedi, A. K., Pasillas, R., et al. (2020). Long-term effect of a nonrandomized psychosocial mindfulness-based intervention in Hispanic/Latina breast cancer survivors. *Integrative Cancer Therapy*, 19, 1534735419890682. <https://doi.org/10.1177/1534735419890682>.
 104. Napoles, A. M., Santoyo-Olsson, J., Chacon, L., Stewart, A. L., Dixit, N., & Ortiz, C. (2019). Feasibility of a mobile phone app and telephone coaching survivorship care planning program among Spanish-speaking breast cancer survivors. *JMIR Cancer*, 5(2), e13543. <https://doi.org/10.2196/13543>.
 105. Yanez, B., Oswald, L. B., Baik, S. H., Buitrago, D., Iacobelli, F., Perez-Tamayo, A., et al. (2020). Brief culturally informed smartphone interventions decrease breast cancer symptom burden among Latina breast cancer survivors. *Psycho-Oncology*, 29(1), 195–203.
 106. Clarke, T. C., Christ, S. L., Soler-Vila, H., Lee, D. J., Arheart, K. L., Prado, G., et al. (2015). Working with cancer: Health and employment among cancer survivors. *Annals of Epidemiology*, 25(11), 832–838. <https://doi.org/10.1016/j.annepidem.2015.07.011>.
 107. Ashing-Giwa, K. T., Lim, J. W., & Gonzalez, P. (2010). Exploring the relationship between physical well-being and healthy lifestyle changes among European- and Latina-American breast and cervical cancer survivors. *Psychooncology*, 19(11), 1161–1170. <https://doi.org/10.1002/pon.1687>.
 108. Fu, O. S., Crew, K. D., Jacobson, J. S., Greenlee, H., Yu, G., Campbell, J., et al. (2009). Ethnicity and persistent symptom burden in breast cancer survivors. *Journal of Cancer Survivorship*, 3(4), 241–250. <https://doi.org/10.1007/s11764-009-0100-7>.
 109. Janz, N. K., Mujahid, M. S., Hawley, S. T., Griggs, J. J., Alderman, A., Hamilton, A. S., et al. (2009). Racial/ethnic differences in quality of life after diagnosis of breast cancer. *Journal of Cancer Survivorship*, 3(4), 212–222. <https://doi.org/10.1007/s11764-009-0097-y>.
 110. Paxton, R. J., Phillips, K. L., Jones, L. A., Chang, S., Taylor, W. C., Courneya, K. S., et al. (2012). Associations among physical activity, body mass index, and health-related quality of life by race/ethnicity in a diverse sample of breast cancer survivors. *Cancer*, 118(16), 4024–4031. <https://doi.org/10.1002/cncr.27389>.
 111. Ross, L. E., Hall, I. J., Fairley, T. L., Taylor, Y. J., & Howard, D. L. (2008). Prayer and self-reported health among cancer survivors in the United States, National Health Interview Survey, 2002. *Journal of Alternative and Complementary Medicine*, 14(8), 931–938. <https://doi.org/10.1089/acm.2007.0788>.
 112. Armbruster, S. D., Song, J., Gatus, L., Lu, K. H., & Basen-Engquist, K. M. (2018). Endometrial cancer survivors' sleep patterns before and after a physical activity intervention: A retrospective cohort analysis. *Gynecological Oncology*, 149(1), 133–139. <https://doi.org/10.1016/j.ygyno.2018.01.028>.
 113. Park, J., Wehrlen, L., Mitchell, S. A., Yang, L., & Bevans, M. F. (2019). Fatigue predicts impaired social adjustment in survivors of allogeneic hematopoietic cell transplantation (HCT). *Supportive Care in Cancer*, 27(4), 1355–1363. <https://doi.org/10.1007/s00520-018-4411-y>.
 114. Zhou, E. S., Clark, K., Recklitis, C. J., Obenchain, R., & Loscalzo, M. (2018). Sleepless from the get go: Sleep problems prior to initiating cancer treatment. *International Journal of Behavioral Medicine*, 25(5), 502–516. <https://doi.org/10.1007/s12529-018-9715-2>.
 115. Bellizzi, K. M., Aziz, N. M., Rowland, J. H., Weaver, K., Arora, N. K., Hamilton, A. S., et al. (2012). Double jeopardy? Age, race, and HRQOL in older adults with cancer. *Journal of Cancer Epidemiology*. <https://doi.org/10.1155/2012/478642>.
 116. Blinder, V., Patil, S., Eberle, C., Griggs, J., & Maly, R. C. (2013). Early predictors of not returning to work in low-income breast cancer survivors: A 5-year longitudinal study. *Breast Cancer Research and Treatment*, 140(2), 407–416. <https://doi.org/10.1007/s10549-013-2625-8>.
 117. Orom, H., Biddle, C., Underwood, W., 3rd, Homish, G. G., & Olsson, C. (2017). Racial/ethnic and socioeconomic disparities in prostate cancer survivors' prostate-specific quality of life. *Urology*. <https://doi.org/10.1016/j.urolgy.2017.08.014>.
 118. Lee, E., Takita, C., Wright, J. L., Reis, I. M., Zhao, W., Nelson, O. L., et al. (2016). Characterization of risk factors for adjuvant radiotherapy-associated pain in a tri-racial/ethnic breast cancer population. *Pain*, 157(5), 1122–1131. <https://doi.org/10.1097/j.pain.0000000000000489>.
 119. Krok-Schoen, J. L., Fernandez, K., Unzeitig, G. W., Rubio, G., Paskett, E. D., Post, D. M., et al. (2019). Hispanic breast cancer patients' symptom experience and patient-physician communication during chemotherapy. *Supportive Care in Cancer*, 27(2), 697–704. <https://doi.org/10.1007/s00520-018-4375-y>.
 120. Meneses, K., Gisiger-Camata, S., Benz, R., Raju, D., Bail, J. R., Benitez, T. J., et al. (2018). Telehealth intervention for Latina breast cancer survivors: A pilot. *Womens Health (London)*, 14, 1745506518778721. <https://doi.org/10.1177/1745506518778721>.
 121. Segrin, C., Badger, T. A., Sikorskii, A., Pasvogel, A., Weihs, K., Lopez, A. M., et al. (2019). Longitudinal dyadic interdependence in psychological distress among Latinas with breast

- cancer and their caregivers. *Supportive Care in Cancer*. <https://doi.org/10.1007/s00520-019-05121-4>.
122. Sleight, A. G., Lyons, K. D., Vigen, C., Macdonald, H., & Clark, F. (2019). The association of health-related quality of life with unmet supportive care needs and sociodemographic factors in low-income Latina breast cancer survivors: A single-Centre pilot study. *Disability and Rehabilitation*, 41(26), 3151–3156.
123. Owens, B., Jackson, M., & Berndt, A. (2009). Complementary therapy used by Hispanic women during treatment for breast cancer. *Journal of Holistic Nursing*, 27(3), 167–176.
124. Black, D. S., Cole, S. W., Christodoulou, G., & Figueiredo, J. C. (2018). Genomic mechanisms of fatigue in survivors of colorectal cancer. *Cancer*, 124(12), 2637–2644. <https://doi.org/10.1002/cncr.31356>.
125. Sanchez-Birkhead, A. C., Carbajal-Salisbury, S., Larreta, J. A., Lovlien, L., Hendricks, H., Dingley, C., et al. (2017). A community-based approach to assessing the physical, emotional, and health status of hispanic breast cancer survivors. *Hispanic Health Care International*, 15(4), 166–172. <https://doi.org/10.1177/1540415317738016>.
126. Brisbois, M. D. (2014). An interpretive description of chemotherapy-induced premature menopause among Latinas with breast cancer. *Oncology Nursing Forum*, 41(5), E282–E289. <https://doi.org/10.1188/14.onf.e282-e289>.
127. Ashing-Giwa, K. T. (2008). Enhancing physical well-being and overall quality of life among underserved Latina-American cervical cancer survivors: Feasibility study. *Journal of Cancer Survivorship*, 2(3), 215–223. <https://doi.org/10.1007/s11764-008-0061-2>.
128. Ashing-Giwa, K. T., & Lim, J. W. (2010). Exploring the association between functional strain and emotional well-being among a population-based sample of breast cancer survivors. *Psychooncology*, 19(2), 150–159. <https://doi.org/10.1002/pon.1517>.
129. Dyer, K. E. (2015). "Surviving is not the same as living": Cancer and Sobrevivencia in Puerto Rico. *Social Science and Medicine*, 132, 20–29. <https://doi.org/10.1016/j.socscimed.2015.02.033>.
130. Ashing-Giwa, K. T., Padilla, G., Tejero, J., Kraemer, J., Wright, K., Coscarelli, A., et al. (2004). Understanding the breast cancer experience of women: A qualitative study of African American, Asian American. *Latina and Caucasian cancer survivors*. *Psychooncology*, 13(6), 408–428. <https://doi.org/10.1002/pon.750>.
131. Castillo, A., Mendiola, J., & Tiemensma, J. (2019). Emotions and coping strategies during breast cancer in Latina women: A focus group study. *Hispanic Health Care International*, 17(3), 96–102. <https://doi.org/10.1177/1540415319837680>.
132. Casillas, J., Kahn, K. L., Doose, M., Landier, W., Bhatia, S., Hernandez, J., et al. (2010). Transitioning childhood cancer survivors to adult-centered healthcare: Insights from parents, adolescent, and young adult survivors. *Psychooncology*, 19(9), 982–990. <https://doi.org/10.1002/pon.1650>.
133. Crookes, D. M., Shelton, R. C., Tehranifar, P., Aycinena, C., Gaffney, A. O., Koch, P., et al. (2016). Social networks and social support for healthy eating among Latina breast cancer survivors: Implications for social and behavioral interventions. *Journal of Cancer Survivorship*, 10(2), 291–301. <https://doi.org/10.1007/s11764-015-0475-6>.
134. Galvan, N., Buki, L. P., & Garces, D. M. (2009). Suddenly, a carriage appears: Social support needs of Latina breast cancer survivors. *Journal of Psychosocial Oncology*, 27(3), 361–382. <https://doi.org/10.1080/07347330902979283>.
135. Kent, E. E., Alfano, C. M., Smith, A. W., Bernstein, L., McTier, A., Baumgartner, K. B., et al. (2013). The roles of support seeking and race/ethnicity in posttraumatic growth among breast cancer survivors. *Journal of Psychosocial Oncology*, 31(4), 393–412. <https://doi.org/10.1080/07347332.2013.798759>.
136. Napoles, A. M., Ortiz, C., Santoyo-Olsson, J., Stewart, A. L., Lee, H. E., Duron, Y., et al. (2017). Post-treatment survivorship care needs of Spanish-speaking Latinas with breast cancer. *Journal of Community Support and Oncology*, 15(1), 20–27. <https://doi.org/10.12788/jcso.0325>.
137. Patel, S. K., Lo, T. T., Dennis, J. M., & Bhatia, S. (2013). Neurocognitive and behavioral outcomes in Latino childhood cancer survivors. *Pediatric Blood & Cancer*, 60(10), 1696–1702. <https://doi.org/10.1002/pbc.24608>.
138. Sammarco, A., & Konecny, L. M. (2010). Quality of life, social support, and uncertainty among Latina and Caucasian breast cancer survivors: A comparative study. *Oncology Nursing Forum*, 37(1), 93–99. <https://doi.org/10.1188/10.onf.93-99>.
139. Smith, A. B., & Bashore, L. (2006). The effect of clinic-based health promotion education on perceived health status and health promotion behaviors of adolescent and young adult cancer survivors. *Journal of Pediatric Oncology Nursing*, 23(6), 326–334.
140. Lockhart, J. S., Oberleitner, M. G., & Nolfi, D. A. (2018). The Hispanic/Latino immigrant cancer survivor experience in the United States: A scoping review. *Annual Review of Nursing Research*, 37(1), 119–160.
141. Moreno, P. I., Ramirez, A. G., San Miguel-Majors, S. L., Castillo, L., Fox, R. S., Gallion, K. J., et al. (2019). Unmet supportive care needs in Hispanic/Latino cancer survivors: Prevalence and associations with patient-provider communication, satisfaction with cancer care, and symptom burden. *Supportive Care in Cancer*, 27(4), 1383–1394. <https://doi.org/10.1007/s00520-018-4426-4>.
142. Moreno, P. I., Ramirez, A. G., San Miguel-Majors, S. L., Fox, R. S., Castillo, L., Gallion, K. J., et al. (2018). Satisfaction with cancer care, self-efficacy, and health-related quality of life in Latino cancer survivors. *Cancer*, 124(8), 1770–1779. <https://doi.org/10.1002/cncr.31263>.
143. Ochoa, C. Y., Haardorfer, R., Escoffery, C., Stein, K., & Alcaraz, K. I. (2018). Examining the role of social support and spirituality on the general health perceptions of Hispanic cancer survivors. *Psychooncology*, 27(9), 2189–2197. <https://doi.org/10.1002/pon.4795>.
144. Segrin, C., Badger, T. A., & Sikorskii, A. (2019). A dyadic analysis of loneliness and health-related quality of life in Latinas with breast cancer and their informal caregivers. *Journal of Psychosocial Oncology*, 37(2), 213–227.
145. Sleight, A. G., Lyons, K. D., Vigen, C., Macdonald, H., & Clark, F. (2018). Supportive care priorities of low-income Latina breast cancer survivors. *Supportive Care in Cancer*, 26(11), 3851–3859.
146. Blinder, V., Eberle, C., Patil, S., Gany, F. M., & Bradley, C. J. (2017). Women with breast cancer who work for accommodating employers more likely to retain jobs after treatment. *Health Affairs (Millwood)*, 36(2), 274–281. <https://doi.org/10.1377/hlthaff.2016.1196>.
147. Blinder, V. S., Patil, S., Thind, A., Diamant, A., Hudis, C. A., Basch, E., et al. (2012). Return to work in low-income Latina and non-Latina white breast cancer survivors: A 3-year longitudinal study. *Cancer*, 118(6), 1664–1674. <https://doi.org/10.1002/cncr.26478>.
148. Burg, M. A., Adorno, G., Lopez, E. D., Loerzel, V., Stein, K., Wallace, C., et al. (2015). Current unmet needs of cancer survivors: Analysis of open-ended responses to the American Cancer Society Study of Cancer Survivors II. *Cancer*, 121(4), 623–630. <https://doi.org/10.1002/cncr.28951>.
149. Jagsi, R., Pottow, J. A., Griffith, K. A., Bradley, C., Hamilton, A. S., Graff, J., et al. (2014). Long-term financial burden of breast cancer: Experiences of a diverse cohort of survivors identified through population-based registries. *Journal of Clinical Oncology*, 32(12), 1269–1276. <https://doi.org/10.1200/jco.2013.53.0956>.

150. Mujahid, M. S., Janz, N. K., Hawley, S. T., Griggs, J. J., Hamilton, A. S., Graff, J., et al. (2011). Racial/ethnic differences in job loss for women with breast cancer. *Journal of Cancer Survivorship*, 5(1), 102–111. <https://doi.org/10.1007/s11764-010-0152-8>.
151. Mujahid, M. S., Janz, N. K., Hawley, S. T., Griggs, J. J., Hamilton, A. S., & Katz, S. J. (2010). The impact of sociodemographic, treatment, and work support on missed work after breast cancer diagnosis. *Breast Cancer Research and Treatment*, 119(1), 213–220. <https://doi.org/10.1007/s10549-009-0389-y>.
152. Pisu, M., Kenzik, K. M., Oster, R. A., Drentea, P., Ashing, K. T., Fouad, M., et al. (2015). Economic hardship of minority and non-minority cancer survivors 1 year after diagnosis: Another long-term effect of cancer? *Cancer*, 121(8), 1257–1264. <https://doi.org/10.1002/cncr.29206>.
153. Sleight, A. G. (2017). Occupational Engagement in Low-Income Latina Breast Cancer Survivors. *Am Journal of Occupational Therapy*, 71(2), 1–8. <https://doi.org/10.5014/ajot.2017.023739>.
154. Chebli, P., Lemus, J., Avila, C., Pena, K., Mariscal, B., Merlos, S., et al. (2019). Multilevel determinants of financial toxicity in breast cancer care: Perspectives of healthcare professionals and Latina survivors. *Supportive Care in Cancer*. <https://doi.org/10.1007/s00520-019-05119-y>.
155. Polek, C., Hardie, T., & Deatrick, J. A. (2019). Breast cancer survivorship experiences of urban Hispanic women. *Journal of Cancer Education*. <https://doi.org/10.1007/s13187-019-01543-0>.
156. Santoyo-Olsson, J., Stewart, A. L., Samayoa, C., Palomino, H., Urias, A., Gonzalez, N., et al. (2019). Translating a stress management intervention for rural Latina breast cancer survivors: The Nuevo Amanecer-II. *PLoS ONE*, 14, 10.
157. Smith, G. L., Lopez-Olivo, M. A., Advani, P. G., Ning, M. S., Geng, Y., Giordano, S. H., et al. (2019). Financial burdens of cancer treatment: A systematic review of risk factors and outcomes. *JNCCN Journal of the National Comprehensive Cancer Network*, 17(10), 1184–1192. <https://doi.org/10.6004/jnccn.2019.7305>.
158. Rush, C. L., Lobo, T., Serrano, A., Blasini, M., Campos, C., & Graves, K. D. (2016). Complementary and alternative medicine use and Latina breast cancer survivors' symptoms and functioning. *Healthcare (Basel)*, 4, 4. <https://doi.org/10.3390/healthcare4040080>.
159. Canada, A. L., Fitchett, G., Murphy, P. E., Stein, K., Portier, K., Crammer, C., et al. (2013). Racial/ethnic differences in spiritual well-being among cancer survivors. *Journal of Behavioral Medicine*, 36(5), 441–453. <https://doi.org/10.1007/s10865-012-9439-8>.
160. Munoz, A. R., Salsman, J. M., Stein, K. D., & Cella, D. (2015). Reference values of the functional assessment of chronic illness therapy-spiritual well-being: A report from the American Cancer Society's studies of cancer survivors. *Cancer*, 121(11), 1838–1844. <https://doi.org/10.1002/cncr.29286>.
161. Levine, E. G., Yoo, G., Aviv, C., Ewing, C., & Au, A. (2007). Ethnicity and spirituality in breast cancer survivors. *Journal of Cancer Survivorship*, 1(3), 212–225. <https://doi.org/10.1007/s11764-007-0024-z>.
162. Prince, P., Mitchell, S. A., Wehrlen, L., Childs, R., Savani, B., Yang, L., et al. (2015). Spiritual well-being in hispanic and non-hispanic survivors of allogeneic hematopoietic stem cell transplantation. *Journal of Psychosocial Oncology*, 33(6), 635–654. <https://doi.org/10.1080/07347332.2015.1082167>.
163. Wildes, K. A., Miller, A. R., de Majors, S. S., & Ramirez, A. G. (2009). The religiosity/spirituality of Latina breast cancer survivors and influence on health-related quality of life. *Psychooncology*, 18(8), 831–840. <https://doi.org/10.1002/pon.1475>.
164. Miller, A. M., Ashing, K. T., Modeste, N. N., Herring, R. P., & Sealy, D. A. (2015). Contextual factors influencing health-related quality of life in African American and Latina breast cancer survivors. *Journal of Cancer Survivorship*, 9(3), 441–449. <https://doi.org/10.1007/s11764-014-0420-0>.
165. Rossi, A., Garber, C. E., Kaur, G., Xue, X., Goldberg, G. L., & Nevadunsky, N. S. (2017). Physical activity-related differences in body mass index and patient-reported quality of life in socioculturally diverse endometrial cancer survivors. *Supportive Care in Cancer*, 25(7), 2169–2177. <https://doi.org/10.1007/s00520-017-3622-y>.
166. Ashing-Giwa, K. T., & Lim, J. W. (2011). Health-related quality of life outcomes among cervical cancer survivors: Examining ethnic and linguistic differences. *Cancer Epidemiology*, 35(2), 194–201. <https://doi.org/10.1016/j.canep.2010.06.006>.
167. Graves, K. D., Jensen, R. E., Canar, J., Perret-Gentil, M., Leventhal, K. G., Gonzalez, F., et al. (2012). Through the lens of culture: Quality of life among Latina breast cancer survivors. *Breast Cancer Research and Treatment*, 136(2), 603–613. <https://doi.org/10.1007/s10549-012-2291-2>.
168. Sammarco, A., & Konecny, L. M. (2008). Quality of life, social support, and uncertainty among Latina breast cancer survivors. *Oncology Nursing Forum*, 35(5), 844–849. <https://doi.org/10.1188/08.onf.844-849>.
169. Smith, A. W., Alfano, C. M., Reeve, B. B., Irwin, M. L., Bernstein, L., Baumgartner, K., et al. (2009). Race/ethnicity, physical activity, and quality of life in breast cancer survivors. *Cancer Epidemiology, Biomarkers & Prevention*, 18(2), 656–663. <https://doi.org/10.1158/1055-9965.epi-08-0352>.
170. Wildes, K. A., Miller, A. R., de Majors, S. S., Otto, P. M., & Ramirez, A. G. (2011). The satisfaction of Latina breast cancer survivors with their healthcare and health-related quality of life. *Journal of Womens Health (Larchmt)*, 20(7), 1065–1074. <https://doi.org/10.1089/jwh.2010.2141>.
171. Ramirez, A. G., Choi, B. Y., Munoz, E., Perez, A., Gallion, K. J., Moreno, P. I., et al. (2019). Assessing the effect of patient navigator assistance for psychosocial support services on health-related quality of life in a randomized clinical trial in Latino breast, prostate, and colorectal cancer survivors. *Cancer*. <https://doi.org/10.1002/cncr.32626>.
172. Anderson, K. O., Green, C. R., & Payne, R. (2009). Racial and ethnic disparities in pain: Causes and consequences of unequal care. *The Journal of Pain*, 10(12), 1187–1204. <https://doi.org/10.1016/j.jpain.2009.10.002>.
173. Balsa, A. I., McGuire, T. G., & Meredith, L. S. (2005). Testing for statistical discrimination in health care. *Health Services Research*, 40(1), 227–252. <https://doi.org/10.1111/j.1475-6773.2005.00351.x>.
174. Burgess, D. J., Gravely, A. A., Nelson, D. B., van Ryn, M., Bair, M. J., Kerns, R. D., et al. (2013). A national study of racial differences in pain screening rates in the VA health care system. *The Clinical Journal of Pain*, 29(2), 118–123. <https://doi.org/10.1097/AJP.0b013e31826a86ae>.
175. McGuire, T. G., & Miranda, J. (2008). New evidence regarding racial and ethnic disparities in mental health: Policy implications. *Health Affairs (Millwood)*, 27(2), 393–403. <https://doi.org/10.1377/hlthaff.27.2.393>.
176. Commission on Cancer. (2019). Cancer Program Standards: Ensuring Patient-Centered Care (2016 Edition). Retrieved May 29, 2020 from <https://www.facs.org/quality-programs/cancer/coc/standards>.
177. Zebrack, B., Kayser, K., Bybee, D., Padgett, L., Sundstrom, L., Jobin, C., et al. (2017). A practice-based evaluation of distress screening protocol adherence and medical service utilization. *Journal of the National Comprehensive Cancer Network: JNCCN*, 15(7), 903–912. <https://doi.org/10.6004/jnccn.2017.0120>.

178. National Comprehensive Cancer Network. (2019). NCCN distress thermometer and problem list for patients. Retrieved May 29, 2019 from <https://www.nccn.org/about/permissions/thermometer.aspx>.
179. Basch, E., Deal, A. M., Kris, M. G., Scher, H. I., Hudis, C. A., Sabbatini, P., et al. (2016). Symptom monitoring with patient-reported outcomes during routine cancer treatment: A randomized controlled trial. *Journal of Clinical Oncology*, 34(6), 557–565. <https://doi.org/10.1200/jco.2015.63.0830>.
180. Kaiser Family Foundation. (2019). Uninsured rates for the nonelderly by race/ethnicity. 2017. Retrieved May 29, 2019 from <https://www.kff.org/uninsured/state-indicator/rate-by-raceethnicity/>.
181. Lopez, M. H., Gonzalez-Barrera, A., & Krogstad, J. M. Latinos and discrimination. In: Pew Research Center: Hispanic Trends. 2018. Retrieved May 29, 2019 from <https://www.pewhispanic.org/2018/10/25/latinos-and-discrimination/>.
182. Soto E. (2019). 5 Conversations to Improve Cancer Care for Older Hispanic Adults. American Society of Clinical Oncology: Cancer.Net. 2018. Retrieved May 29, 2019 from <https://www.cancer.net/blog/2018-08/5-conversations-improve-cancer-care-older-hispanic-adults>.
183. Montealegre, J. R., Zhou, R., Amirian, E. S., Follen, M., & Scheurer, M. E. (2013). Nativity disparities in late-stage diagnosis and cause-specific survival among Hispanic women with invasive cervical cancer: An analysis of surveillance, epidemiology, and end results data. *Cancer Causes and Control*, 24(11), 1985–1994. <https://doi.org/10.1007/s10552-013-0274-1>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.